

## Skinvive Post Care Preview



## Skinvive Post Care

***SKINVIVE™ by JUVÉDERM® Injectable Gel Post care***

1. Treatment with SKINVIVE™ by JUVÉDERM® has minimal downtime, especially compared with other skin quality treatment recovery times that can stretch up to 2 weeks.
2. Avoid touching the treated area and keep hair (and phone) away from the treated area for several hours
3. Avoid makeup for 24 hours after treatment
4. Postpone dental procedures for atleast 4 weeks after treatment
5. Avoid using skincare products, extensive sun or heat exposure, strenous exercise, and alcoholic beverages for 24 hours after treatment
6. Bruising, firmness and tenderness around the injection site is normal and expected

Immediately after the treatment, there may be slight redness, swelling, tenderness and an itching sensation in the treated area. This is a normal result of the injection. The inconvenience is usually temporary and generally disappears in less than 6 hours after injection into the skin.

Occasionally some patients experience swelling for about a week. This means that the result directly after the treatment should not be seen as the final result. If you have previously suffered from facial cold sores, there is a risk that the needle punctures could contribute to another eruption of cold sores. If you are using aspirin or any similar medication, be aware that these may increase the bruising and bleeding at the injection site.

Other types of reactions are very rare, but occasionally some patients have experienced other reactions. Please be sure to review the consent form for further details.

These aesthetic products are temporary injectable fillers and the duration of the effect can vary from patient to patient. You may wish to consider additional treatments before the effect has completely dissipated.

[Save Record](#)